arXiv:1610.09658v1 [math.GT] 30 Oct 2016

On moduli spaces of positive scalar curvature metrics on certain manifolds

Michael Wiemeler*

We show that the homotopy groups of moduli spaces of positive scalar curvature metrics on certain simply connected spin manifolds are non-trivial.

1 Introduction

In recent years a lot of work was devoted on the study of spaces $\mathcal{R}^+(M)$ of Riemannian metrics of positive scalar curvature on a given spin manifold M. Here we only want to mention the works [BERW14] and [HSS14].

The first work relates the study of the homotopy type of these spaces to certain infinite loop spaces. It is shown in this paper that infinitely many homotopy groups of these spaces are non-trivial.

In the second work it was shown that in degrees d much smaller than the dimension n of the manifold M there are classes in the the homotopy groups $\pi_d(\mathcal{R}^+(M))$, which descent to non-trivial classes in the homotopy groups of the observer moduli space of positive scalar curvature metrics on M, if M fulfills some technical condition.

In this note we combine the results and methods of the above mentioned papers to prove the following theorem.

Theorem 1.1 *Let M be a closed spin manifold of dimension $n \geq 6$ which admits a metric of positive scalar curvature. If $k = 4s - n - 1 > 2n$, $s \in \mathbb{Z}$ and $g_0 \in \mathcal{R}^+(M)$, then*

$$\pi_k(\mathcal{R}^+(M), g_0) \otimes \mathbb{Q} \to \pi_k(\mathcal{M}^+_{observer}(M), g_0) \otimes \mathbb{Q}$$

has non-trivial image.

If moreover, there is no non-trivial orientation-preserving action of a finite group on M, then

$$\pi_k(\mathcal{R}^+(M), g_0) \otimes \mathbb{Q} \to \pi_k(\mathcal{M}^+_0(M), g_0) \otimes \mathbb{Q}$$

has non-trivial image.

*The research for this work was supported by DFG-Grant HA 3160/6-1.

Here $\mathcal{M}_0^+(M) = \mathcal{R}^+(M)/\operatorname{Diff}_0(M)$ denotes the moduli space of metrics of positive scalar curvature with respect to the group $\operatorname{Diff}_0(M)$ of orientation-preserving diffeomorphisms of M.

Examples of manifolds on which no finite group acts non-trivially and orientation-preserving have been given by Puppe [Pup95]. These examples are simply connected spin manifolds of dimension six. Therefore they admit metrics of positive scalar curvature.

Moreover, we show the following result:

Theorem 1.2 *Let $n > 1$. Then in the image of the map*

$$\pi_1(\mathcal{R}^+(S^{4n-2})) \to \pi_1(\mathcal{M}_0^+(S^{4n-2}))$$

there are elements of infinite order.

As far as we know, these are the first examples of elements in the fundamental group of the space of positive scalar curvature metrics which decent to elements of infinite order in the fundamental group of $\mathcal{M}_0^+(S^{4n-2})$. Note that, by [Wal14], $\pi_1(\mathcal{R}^+(S^{4n-2}))$ is abelian. The space $\mathcal{R}^+(S^2)$ is known to be contractible [RS01]. Hence, the above theorem is false for $n = 1$.

This note is structured as follows. In the next Section 2 we proof Theorem 1.1. In Section 3 we recall some transversality results in the context of infinite dimensional manifolds. Then in Section 4 we recall Ebin's slice theorem and some of its consequences. In the Section 5 we give the proof of Theorem 1.2. Finally in Section 6 we show that our methods are not fine enough to deal with the moduli space of positive scalar curvature metrics with respect to the full diffeomorphism group.

2 The proof of Theorem 1.1

Our proof of Theorem 1.1 is based on the following result:

Theorem 2.1 ([BERW14, Theorem A]) *Let M be a closed spin manifold of dimension $n \geq 6$. If $k = 4s - n - 1 \geq 0$ and $g_0 \in \mathcal{R}^+(M)$, then the map*

$$A_k \otimes \mathbb{Q} : \pi_k(\mathcal{R}^+(M), g_0) \otimes \mathbb{Q} \to \mathbb{Q}$$

is surjective.

Here A_k denotes the secondary index invariant for metrics of positive scalar curvature metrics. There are two definitions for this invariant one is by Hitchin [Hit74], the other is by Gromov and Lawson [GL83].

As shown in [Ebe14], these two definitions lead to the same invariant. This is important for our work because we use results of [BERW14], where mainly the first definition is used, and of [HSS14], where the definition of Gromov and Lawson is used.

We also need the following definition from [HSS14]:

Definition 2.2 *Let M be an oriented closed smooth manifold. We call M a $\hat{A}$-multiplicative fiber in degree k if for every oriented fiber bundle $M \to E \to S^{k+1}$ we have $\hat{A}(E) = 0$.*

For $\hat{A}$-multiplicative manifolds the following is known:

Theorem 2.3 *Let M be a closed spin manifold and $k \geq 2$. If M is a $\hat{A}$-multiplicative fiber in degree k, then the following holds:*

1. *The map $A_k \otimes \mathbb{Q}$ from above factors through $\pi_k(\mathcal{M}^+_{observer}(M), g_0) \otimes \mathbb{Q}$.*
2. *If there is no non-trivial smooth orientation-preserving action of a finite group on M then $A_k \otimes \mathbb{Q}$ factors through $\pi_k(\mathcal{M}^+_0(M), g_0) \otimes \mathbb{Q}$.*

Proof. The first statement is proved in [HSS14, Section 2]. The proof of the second statement is similar. Since there is no non-trivial orientation-preserving smooth action of a finite group on M the isometry group of any Riemannian metric on M contains at most one orientation-reversing involution and the identity. Therefore the group $\mathrm{Diff}_0(M)$ of orientation-preserving diffeomorphisms acts freely on $\mathcal{R}^+(M)$.

In particular, there is an exact sequence

$$\pi_k(\mathrm{Diff}_0(M)) \otimes \mathbb{Q} \to \pi_k(\mathcal{R}^+(M)) \otimes \mathbb{Q} \to \pi_k(\mathcal{M}^+_0) \otimes \mathbb{Q}$$

Since M is an $\hat{A}$-multiplicative fiber in degree k it follows from the arguments in [HSS14, Section 2] that $A_k \otimes \mathbb{Q}$ vanishes on the image of $\pi_k(\mathrm{Diff}_0(M)) \otimes \mathbb{Q}$. Therefore the theorem follows. □

Our Theorem 1.1 now follows from the above two theorems and the following lemma.

Lemma 2.4 *Let M be a closed oriented smooth manifold of dimension $n \geq 1$. If $k > 2n$, then M is a $\hat{A}$-multiplicative fiber in degree k.*

Proof. Let $M \to E \to S^{k+1}$ be a smooth oriented fiber bundle. The tangent bundle of E is isomorphic to $p^*(TS^{k+1}) \oplus V$ where V is the bundle along the fiber. Since TS^{k+1} is stably trivial, it follows that TE and V are stably isomorphic. Therefore the Pontrjagin classes of E are concentrated in degrees smaller or equal to $2n$. Moreover, since $n < k$, it follows from an inspection of the Serre spectral sequence for the fibration that $H^j(E; \mathbb{Q}) = 0$ for $n < j < k + 1$. Because $2n < k$ the Pontrjagin classes of E are concentrated in degrees smaller or equal to n. Moreover it follows that all products of these classes of degree greater than n must vanish. In particular all the Pontrjagin numbers of E vanish and the theorem is proved. □

3 Transversality

In this section we start to collect the necessary material for the proof of Theorem 1.2. We need the following transversality result for maps into infinite dimensional manifolds.

Lemma 3.1 *Let F be a topological vector space and $p : F \to \mathbb{R}^{n+1}$ a continuous linear surjective map and $U \subset F$ an open neighborhood of zero.*

If $f : D^n \to F$ is a continuous map and $K_1, \dots, K_m \subset D^n$ are compact such that there are open $U_1, \dots, U_m \subset U$ with $f(K_i) \subset U_i$, then there is a continuous map $f' : D^n \to U$, such that

1. *$f'(D^n_{1/2}) \subset U - \{0\}$,*
2. *there is a homotopy h_t from f to f', such that $h_t(K_i) \subset U_i$ for all t and i and $h_t|_{\partial D^n} = f|_{\partial D^n}$.*

Proof. The existence of p guarantees that F is isomorphic as a topological vector space to $\mathbb{R}^{n+1} \times F'$ for some closed subvector space $F' \subset F$. Therefore the Lemma follows from the finite dimensional case $F = \mathbb{R}^{n+1}$. This case follows from transversality considerations. □

Remark 3.2 *If there is a continuous scalar product on an infinite dimensional topological vector space, then there is always a map p as above. One can take the orthogonal projection onto some $n+1$-dimensional subvector space.*

4 Consequences of Ebin's slice theorem

Ebin [Ebi70] showed the following theorem:

Theorem 4.1 *Let $\mathcal{R}$ be the space of Riemannian metrics on a closed manifold M, $g \in \mathcal{R}$.*

Then there is a neighborhood N of g, which is homeomorphic to $V \times D(E)$, where V is a neighborhood of $\mathrm{Id} \in \mathrm{Diff}(M)/\mathrm{Iso}(g)$ and $D(E)$ is an invariant contractible open subset containing 0 in some infinite dimensional $\mathrm{Iso}(g)$-representation E.

The homeomorphism $\phi : V \times D(E) \to N$ in the above theorem is of the following form. There is a section $\chi : V \to \mathrm{Diff}(M)$ to the map $\mathrm{Diff}(M) \to \mathrm{Diff}(M)/\mathrm{Iso}(g)$ and the homeomorphism satisfies $\phi(v, e) = \chi(v)\phi(\mathrm{Id}, e)$.

Here the isometry group $\mathrm{Iso}(g)$ of g is a compact Lie subgroup of $\mathrm{Diff}(M)$, since M is compact.

From the proof of this theorem it follows that E is isomorphic as a topological vector space to $E^{\mathrm{Iso}(g)} \times F$, where F is a $\mathrm{Iso}(g)$-invariant closed subspace of E. Moreover, there is a continuous scalar product on E so that the remark of the previous section applies

to E and F. Note that Bourguignon [Bou75, Proposition III.20] has shown that F is infinite dimensional if $\mathrm{Iso}(g)$ is non-trivial.

For $g' \in N$, $\mathrm{Iso}(g')$ is conjugated in $\mathrm{Diff}(M)$ to a subgroup of $\mathrm{Iso}(g)$ (see [Ebi70, Theorem 8.1]). Furthermore, it follows from the proof of that theorem that the minimal stratum in N, i.e. the set of those metrics in N whose isometry group is conjugated to $\mathrm{Iso}(g)$, is given by $V \times D(E)^{\mathrm{Iso}(g)}$. In particular, maps from finite dimensional manifolds to N can be made transverse to the minimal stratum.

If $\gamma : I \to N/\mathrm{Diff}(M) = D(E)/\mathrm{Iso}(g)$ is a path, then this path can be lifted to a path in N (see [Bre72, Chapter II.6]).

Denote by $\mathcal{M}$ the moduli space of all Riemannian metrics on the closed manifold M and by $\mathcal{M}_{m,k}$ the moduli space of Riemannian metrics on M with $\dim \mathrm{Iso}(g) < m$ or $(\dim \mathrm{Iso}(g) = m$ and $|\mathrm{Iso}(g)/\mathrm{Iso}(g)^0| \leq k)$.

Then we have:

1. $\mathcal{M}$ is locally homeomorphic to orbit spaces of infinite dimensional representations of compact Lie groups. We call the charts which correspond to these local models "Ebin charts", denoted by E/G.

2. $\mathcal{M}_{m,k-1} \subset \mathcal{M}_{m,k} \subset \mathcal{M}$ are inclusions of open set because each $g \in \mathcal{M}_{m,k-1}$ has a neighborhood N' such that for each $g' \in N'$, $\mathrm{Iso}(g')$ is conjugated to a subgroup of $\mathrm{Iso}(g)$.

Since S^n and D^{n+1} are compact for all $n \in \mathbb{N}$ the following statements follow from the second point above.

3. $\pi_n(\mathcal{M}_{m,0}) = \lim_k \pi_n(\mathcal{M}_{m-1,k})$

4. $\pi_n(\mathcal{M}) = \lim_m \pi_n(\mathcal{M}_{m,0})$

5. $\ker(\pi_n(\mathcal{M}_{m,k}) \to \pi_n(\mathcal{M}_{m+1,0})) = \lim_j \ker(\pi_n(\mathcal{M}_{m,k}) \to \pi_n(\mathcal{M}_{m,j}))$

6. $\ker(\pi_n(\mathcal{M}_{m,0}) \to \pi_n(\mathcal{M})) = \lim_j \ker(\pi_n(\mathcal{M}_{m,0}) \to \pi_n(\mathcal{M}_{j,0}))$

7. similar statements hold for homology groups.

5 The proof of Theorem 1.2

In the following $\mathcal{R}$, $\mathcal{M}$, etc. can be defined as in the previous section, or as $\mathrm{Diff}(M)$-invariant open subspaces of these spaces.

Lemma 5.1 *The map $\pi_1(\mathcal{M}_{m,k-1}) \to \pi_1(\mathcal{M}_{m,k})$ is surjective.*

Proof. Let $\gamma : I \to \mathcal{M}_{m,k}$ with $\gamma(0) = \gamma(1) \in \mathcal{M}_{m,k-1}$.

Then there are finitely many intervals $[a_j, b_j] \subset]0,1[$, so that $\gamma([a_j, b_j])$ is contained in an Ebin chart and $\gamma^{-1}(\mathcal{M}_{m,k} - \mathcal{M}_{m,k-1}) \subset \bigcup_j [a_j, b_j]$.

The path $\gamma|_{[a_j,b_j]}$ can be lifted to a path in an Ebin slice E. There it can be made transversal to the minimal stratum. Since the complement of the minimal stratum in an Ebin chart is connected, γ is homotopic to a closed curve in $\mathcal{M}_{m,k-1}$. □

Lemma 5.2 *The kernel of the map $\pi_1(\mathcal{M}_{m,k-1}) \to \pi_1(\mathcal{M}_{m,k})$ is generated by torsion elements.*

Proof. Let $\gamma : I \to \mathcal{M}_{m,k-1}$ be a closed curve and $\lambda : I \times I \to \mathcal{M}_{m,k}$ a null homotopy of γ.

Then there are finitely many discs $D_1^2, \dots, D_j^2 \subset I \times I - \partial(I \times I)$ with piecewise C^1-boundary, such that:

1. $\lambda^{-1}(\mathcal{M}_{m,k} - \mathcal{M}_{m,k-1}) \subset \bigcup_i (D_i^2 - \partial D_i^2)$
2. $\lambda(D_i^2)$ is contained in an Ebin chart.

Without loss of generality we may assume, that there is a curve $\sigma : I \to I \times I$ from the base point to the boundary of D_k^2 such that $\sigma(t) \notin \bigcup D_i^2$ for $t \neq 1$.

Cutting $I \times I$ along σ leads to a homotopy from γ to $\lambda(\sigma * \partial D_k^2 * \sigma^{-1})$.

Now we can lift $\lambda(\partial D_k^2)$ to a curve in an Ebin slice and make it transversal to the minimal stratum. This leads to a homotopy $\lambda' : I \times I \to \mathcal{M}_{m,k}$ from γ to $\lambda(\sigma * \partial (D_k^2)' * \sigma^{-1})$ with

1. $\lambda'(\partial(I \times I)) \subset \mathcal{M}_{m,k-1}$
2. $\lambda'^{-1}(\mathcal{M}_{m,k} - \mathcal{M}_{m,k-1})$ is contained in $k-1$ discs as above.

By induction we see that γ is homotopic in $\mathcal{M}_{m,k-1}$ to

$$\sigma_1 * \partial D_1^2 * \sigma_1^{-1} * \sigma_2 * \partial D_2^2 \cdots * \sigma_k^{-1}.$$

Therefore it suffices to show that closed curves δ which are contained in an Ebin chart are torsion elements.

To do so, lift δ to an Ebin slice. Let δ' be the lift of δ. Let G be the compact Lie group which acts on this slice. Then there is a $g \in G$, such that $g\delta'(0) = \delta'(1)$. Let k be the order of the class of g in G/G^0.

Then

$$\delta' * g\delta' * \cdots * g^{k-1}\delta'$$

is a lift of δ^k, which ends and starts in the same component of $G\delta'(0)$. Therefore δ^k can be lifted to a closed curve in the Ebin slice. Since the complement of the minimal stratum in the Ebin slice is contractible, it follows that δ is torsion. □

Corollary 5.3 *The kernel of $H_1(\mathcal{M}_{0,1}) \to H_1(\mathcal{M})$ is torsion.*

Proof. It follows from the two lemmas above that the kernels of the maps $H_1(\mathcal{M}_{m,k-1}) \to H_1(\mathcal{M}_{m,k})$ are torsion.

To see this note that $H_1(\mathcal{M}_{m,k}) = \pi_1(\mathcal{M}_{m,k})/C_{m,k}$, where $C_{m,k}$ denotes the commutator subgroup.

Let $\gamma \in \pi_1(\mathcal{M}_{m,k-1})$ such that $[\gamma] = 0 \in H_1(\mathcal{M}_{m,k})$. We have to show that γ is contained in a normal subgroup of $\pi_1(\mathcal{M}_{m,k-1})$ which is generated by torsion elements and $C_{m,k-1}$. Since $[\gamma] = 0 \in H_1(\mathcal{M}_{m,k})$, we have $\gamma \in C_{m,k} \subset \pi_1(\mathcal{M}_{m,k})$. Because $C_{m,k-1} \to C_{m,k}$ is surjective by the first lemma, it follows from the second lemma that up to torsion elements γ is contained in $C_{m,k-1}$. This proves the claim.

The next step is to show that the kernels of $H_1(\mathcal{M}_{0,1}) \to H_1(\mathcal{M}_{1,0})$ and $H_1(\mathcal{M}_{m-1,0}) \to H_1(\mathcal{M}_{m,0})$ are torsion. This follows from 5 and 7 in the previous section, by taking the limit.

Therefore the statement follows from 6 and 7 in the previous section, by taking the limit. □

Corollary 5.4 *Let $n > 1$. Then in the image of the map*

$$\pi_1(\mathcal{R}^+(S^{4n-2})) \to \pi_1(\mathcal{M}_0^+(S^{4n-2}))$$

there are elements of infinite order.

Proof. It follows from transversality considerations that $\pi_1(\mathcal{R}^+_{0,1}) \to \pi_1(\mathcal{R}^+)$ is an isomorphism. Using Theorem 2.1 one sees as in the proof of Theorem 2.3 that there are elements of infinite order in the image of the map $\pi_1(\mathcal{R}^+_{0,1}(S^{4n-2})) \to \pi_1(\mathcal{M}^+_{0,1}(S^{4n-2}))$, whose image in H_1 has infinite order. Note here that spheres are $\hat{A}$-multiplicative fibers in any degree by [HSS14, Proposition 1.9]. Therefore the statement follows from the above corollary. □

6 Remarks on orientation-reversing isometries

In this section, we show that the index difference $A_k \otimes \mathbb{Q}$ cannot be used to detect non-trivial elements in $\pi_k(\mathcal{M}^+(M)) \otimes \mathbb{Q}$ if there is an orientation-reversing diffeomorphism φ of finite order in $\mathrm{Diff}(M)$ which leaves some metric g_0 of positive scalar curvature invariant. Here $\mathcal{M}^+(M)$ denotes the moduli space of positive scalar curvature metrics on the spin manifold M with respect to the full diffeomorphism group of M.

To be more precise we have the following theorem.

Theorem 6.1 *Let M, φ and g_0 as above. Assume $H^1(M;\mathbb{Z}_2) = 0$. Then any element in the image of the map $\pi_k(\mathcal{R}^+, g_0) \otimes \mathbb{Q} \to \pi_k(\mathcal{M}^+, g_0) \otimes \mathbb{Q}$ can be represented by an element $g' \in \pi_k(\mathcal{R}^+, g_0) \otimes \mathbb{Q}$ with $A_k(g') = 0$, $k \geq 2$.*

Proof. Let m be the order of φ and ζ a primitive m-th root of unity. We first prove the theorem with $\mathbb{Q}$ replaced by $\mathbb{Q}[\zeta]$ and will then indicate how the theorem follows.

Since g_0 is fixed by the action of φ on $\mathcal{R}^+$, there is a natural induced action of φ on $\pi_k(\mathcal{R}^+, g_0) \otimes \mathbb{Q}[\zeta]$. This $\mathbb{Q}[\zeta]$-vector space decomposes as a direct sum of the eigenspaces of φ_*.

To see this note that, for $v \in \pi_k(\mathcal{R}^+, g_0) \otimes \mathbb{Q}[\zeta]$,

$$W_v = \langle \varphi_*^l(v);\ l \in \mathbb{Z} \rangle$$

is a finite dimensional φ_*-invariant subvector space of $\pi_k(\mathcal{R}^+, g_0) \otimes \mathbb{Q}[\zeta]$ because φ has finite order. Since the minimal polynomial of the restriction of φ_* to W_v (a divisor of $X^m - 1$), splits over $\mathbb{Q}[\zeta]$ as a product of linear factors each appearing with multiplicity one, $\varphi_*|_{W_v}$ is diagonizable. Because $v \in W_v$, it follows that v is the sum of eigenvectors of φ_*.

The kernel of the natural map $\pi_k(\mathcal{R}^+, g_0) \otimes \mathbb{Q}[\zeta] \to \pi_k(\mathcal{M}^+, g_0) \otimes \mathbb{Q}[\zeta]$ contains all eigenspaces of φ_* except the one corresponding to the eigenvalue 1. This is because, for $g' \in \pi_k(\mathcal{R}^+) \otimes \mathbb{Q}[\zeta]$, $g' - \varphi_* g'$ is clearly contained in this kernel.

Therefore it suffices to show that $A_k \otimes \mathbb{Q}[\zeta]$ vanishes on this eigenspace. To see this we note that for $g' \in \pi_k(\mathcal{R}^+, g_0)$ we have

$$A_k(g') = -A_k(\varphi_* g') \tag{6.1}$$

by the Gromov-Lawson definition of the index difference and the fact that φ is an orientation-reversing isometry of g_0. Note here that since $H^1(M; \mathbb{Z}_2) = 0$ there is a unique spin structure on M. Therefore $\varphi : M \to -M$ is spin preserving.

To get the statement of the theorem with coefficients in $\mathbb{Q}$ and not in $\mathbb{Q}[\zeta]$, note that the Galois group H of the field extension $\mathbb{Q}[\zeta]|\mathbb{Q}$ is finite and acts with fixed point set $\pi_k(\mathcal{R}^+, g_0) \otimes \mathbb{Q}$ on $\pi_k(\mathcal{R}^+, g_0) \otimes \mathbb{Q}[\zeta]$. Therefore, if $g' \in \pi_k(\mathcal{R}^+, g_0) \otimes \mathbb{Q}[\zeta]$ with $A_k(g') = 0$ and $[g'] \in \pi_k(\mathcal{M}^+, g_0) \otimes \mathbb{Q}$, we have

$$A_k\Big(\frac{1}{|H|} \sum_{h \in H} h(g')\Big) = \frac{1}{|H|} \sum_{h \in H} h(A_k(g')) = 0,$$

$$\frac{1}{|H|} \sum_{h \in H} h(g') \in \pi_k(\mathcal{R}^+, g_0) \otimes \mathbb{Q}$$

and

$$\Big[\frac{1}{|H|} \sum_{h \in H} h(g')\Big] = \frac{1}{|H|} \sum_{h \in H} h([g']) = [g'] \in \pi_k(\mathcal{M}^+, g_0) \otimes \mathbb{Q}.$$

Hence, the theorem is proved. □

The examples of Puppe mentioned in the introduction have orientation-reversing involutions. Therefore with our methods one can not do better than in our Theorem 1.1. But we also want to mention that many positively curved manifolds such as spheres and complex projective spaces of odd complex dimensions admit orientation-reversing isometric involutions.

References

[BERW14] Boris Botvinnik, Johannes Ebert, and Oscar Randal-Williams. Infinite loop spaces and positive scalar curvature. Preprint, 2014.

[Bou75] Jean-Pierre Bourguignon. Une stratification de l'espace des structures riemanniennes. *Compositio Math.*, 30:1–41, 1975.

[Bre72] Glen E. Bredon. *Introduction to compact transformation groups*. Academic Press, New York-London, 1972. Pure and Applied Mathematics, Vol. 46.

[Ebe14] Johannes Ebert. The two definitions of the index difference. Preprint, 2014.

[Ebi70] David G. Ebin. The manifold of Riemannian metrics. In *Global Analysis (Proc. Sympos. Pure Math., Vol. XV, Berkeley, Calif., 1968)*, pages 11–40. Amer. Math. Soc., Providence, R.I., 1970.

[GL83] Mikhael Gromov and H. Blaine Lawson, Jr. Positive scalar curvature and the Dirac operator on complete Riemannian manifolds. *Inst. Hautes Études Sci. Publ. Math.*, (58):83–196 (1984), 1983.

[Hit74] Nigel Hitchin. Harmonic spinors. *Advances in Math.*, 14:1–55, 1974.

[HSS14] Bernhard Hanke, Thomas Schick, and Wolfgang Steimle. The space of metrics of positive scalar curvature. *Publ. Math. Inst. Hautes Études Sci.*, 120:335–367, 2014.

[Pup95] Volker Puppe. Simply connected 6-dimensional manifolds with little symmetry and algebras with small tangent space. In *Prospects in topology (Princeton, NJ, 1994)*, volume 138 of *Ann. of Math. Stud.*, pages 283–302. Princeton Univ. Press, Princeton, NJ, 1995.

[RS01] Jonathan Rosenberg and Stephan Stolz. Metrics of positive scalar curvature and connections with surgery. In *Surveys on surgery theory, Vol. 2*, volume 149 of *Ann. of Math. Stud.*, pages 353–386. Princeton Univ. Press, Princeton, NJ, 2001.

[Wal14] Mark Walsh. H-spaces, loop spaces and the space of positive scalar curvature metrics on the sphere. *Geom. Topol.*, 18(4):2189–2243, 2014.